

2,4-Diorganyl 1,3,2,4-dithiadiphosphetane-2-4-disulfides, their structure and S-silyl dithiophosphonic derivatives

Nizamov I., Terenzhev D., Shumatbaev G., Islamov D., Kataeva O., Cherkasov R.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016 Taylor & Francis Group, LLC. Novel 2,4-diorganyl 1,3,2,4-dithiadiphosphetane-2,4-disulfides were prepared by the reactions of tetraphosphorus decasulfide with butylphenyl ether or 3-bromophenyl-1-isoamyl ether in 1,2-dichlorobenzene. Their structures were established by X-ray single crystal diffraction. Chiral S-trimethylsilyl dithiophosphonates were prepared by the reactions of 2,4-diorganyl 1,3,2,4-dithiadiphosphetane-2,4-disulfides with trimethyl silyl ethers of monoterpenols. Unsaturated S-trimethylsilyl dithiophosphonates were synthesized by the reactions of 2,4-diorganyl 1,3,2,4-dithiadiphosphetane-2,4-disulfides with O-trimethylsilyl ethers of geraniol, nerol, and (R,S)-linalool. S-Esters of dithiophosphonic acids, which possessed antimicrobial activity, were prepared by the reactions of chiral S-trimethylsilyl dithiophosphonates with orthoformates, acetals and thioacetals.

<http://dx.doi.org/10.1080/10426507.2016.1216417>

Keywords

1,3,2,4-Dithiadiphosphetane-2,4-disulfides, dithiophosphorylation, monoterpenols, silylation